

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

STUDIES IN MICROBIOLOGY VOLUME 2

Principles of Gene Manipulation

AN INTRODUCTION TO
GENETIC ENGINEERING

R. W. OLD MA, PhD
Department of Biological Sciences,
University of Warwick,
Coventry

S. B. PRIMROSE BSc, PhD
Amersham International plc,
Amersham,
Buckinghamshire

FOURTH EDITION

OXFORD

Blackwell Scientific Publications

LONDON EDINBURGH BOSTON

MELBOURNE PARIS BERLIN VIENNA

© 1980, 1981, 1985, 1989 by
Blackwell Scientific Publications
Editorial offices:
Osney Mead, Oxford OX2 0EL
25 John Street, London WC1N 2BL
23 Ainslie Place, Edinburgh EH3 6AJ
3 Cambridge Center, Cambridge,
Massachusetts 02142, USA
54 University Street, Carlton
Victoria 3053, Australia

Other Editorial Offices:
Arnette SA
2, rue Casimir-Delavigne
75006 Paris
France

Blackwell Wissenschafts-Verlag
Meinekestrasse 4
D-1000 Berlin 15
Germany

Blackwell MZV
Feldgasse 13
A-1238 Wien
Austria

All rights reserved. No part of this
publication may be reproduced, stored
in a retrieval system, or transmitted, in
any form or by any means, electronic,
mechanical, photocopying, recording or
otherwise without the prior permission
of the copyright owner.

First published 1980
Second edition 1981
Reprinted 1982, 1983 (twice)
Third edition 1985
Reprinted 1986 (twice), 1987, 1988
Fourth edition 1989
Reprinted 1991, 1992

Set by Setrite Typesetters, Hong Kong
Printed and bound in Spain by
Printek, S.A., Bilbao

DISTRIBUTORS
Marston Book Services Ltd
PO Box 87
Oxford OX2 0DT
(Orders: Tel: 0865 791155
Fax: 0865 791927
Telex: 837515)

USA
Blackwell Scientific Publications, Inc.
3 Cambridge Center
Cambridge, MA 02142
(Orders: Tel: (800) 759-6102
(617) 225-0401)

Canada
Times Mirror Professional Publishing, Ltd
5240 Finch Avenue East
Scarborough, Ontario M1S 5A2
(Orders: Tel: (800) 268-4178
(416) 298-1588)

Australia
Blackwell Scientific Publications
(Australia) Pty Ltd
54 University Street
Carlton, Victoria 3053
(Orders: Tel: (03) 347-0300)

British Library
Cataloguing in Publication Data
Old, R. W.
Principles of gene manipulation. — 4th ed.
1. Genetic engineering
I. Title II. Series
575.1
ISBN 0-632-02608-1

Library of Congress
Cataloging in Publication Data
Old, R. W.
Principles of gene manipulation: an
introduction to genetic engineering/
R. W. Old, S. B. Primrose. — 4th ed.
p. cm. — (Studies in microbiology; v. 2)
Includes bibliographical references.
ISBN 0-632-02608-1:
1. Genetic engineering. I. Primrose, S.
B. II. Title. III. Series: Studies in
microbiology (Oxford, England); v. 2.
QH442.042 1989
660'.65—dc20




Fig. 4.10 Life cycle and DNA replication of phage M13.




Fig. 5.7 *In-vitro* strand selection of mutants. The DNA-containing sulphur nucleotides is shown in colour.




Fig. 5.9 Procedure for random mutagenesis of a target sequence. In the absence of one of the dNTPs the AMV reverse transcriptase copies the template DNA slowly and inserts any of the dNTPs present for the missing dNTP. On adding back the missing dNTP, replication occurs at its usual high rate. Dots represent mutated bases.

could correct the *CYC-1* mutation, and revertants were obtained that had the expected base sequence. For this method to succeed it is necessary to have stable mutants and a positive selection method for revertants. High DNA concentrations (100 μ g) also are required.

Duplex DNA

Single strands

Single strand +
random primers

Primer extension

Template single
strand

Labelled probe

Denature

Add random-sequence
primersAdd Klenow polymerase
unlabelled dNTPs
labelled dCTP(*)Denature to
release labelled probe

Add to hybridization mix

Fig. 15.2 The random primer method for preparing labelled DNA.




Fig. 15.4 Preparation of probes using a single-stranded m1CE vector.

target and the probe nucleic acid are free to move, thus maximizing the chance that complementary sequences will align and bind. Consequently solution hybridizations go to completion five- to 10-fold faster than those on solid supports (Bryan *et al.* 1986). This can be particularly important in many diagnostic microbiology applications where the concentration of the target sequence is very low and speed is essential.

At the end of the hybridization step it is essential to separate duplexes from unbound probe. If one of the two sequences in the hybridization reaction has been immobilized this separation step is achieved by a simple washing procedure. This explains the popularity of filter hybridization, of which there have been numerous examples in previous chapters.

A variation of the filter hybridization reaction is to attach the probe to the bottom of a microtitre plate well or to a tube (Polsky-Cynkin *et al.* 1985). This facilitates the washing step, reduces the total volume of the hybridization, and facilitates the automated reading of results if a colorimetric detection system is used. A clever utilization of this format is the sandwich hybridization reaction (Ranki *et al.* 1983, Palva & Ranki 1985). Here one probe is attached to the solid support and serves to capture homologous nucleic acids. A second DNA probe, which recognizes a contiguous sequence carries the reporter molecule (Fig. 15.12). Although the sandwich format is far less dependent upon the sample composition than direct blotting methods, it is a relatively slow process. Also, the capture is inefficient, since after denaturing the target DNA, the rate of reassociation of the target with itself in solution is considerably greater than its rate of association with the solid-phase probe (Syvanen *et al.* 1986). For the latter reason RNA probes are preferable to DNA probes.

If solution hybridization has been used, removal of unbound probe is not easy. One method is to digest the single-stranded nucleic acid that remains after hybridization with an appropriate nuclease. Although attractive in principle, in practice efficient digestion and separation are difficult to achieve reliably, particularly with crude samples. An alternative method is to separate duplexes from single-stranded by means of differential binding to hydroxyapatite or antibodies specific for double-stranded nucleic acids. A different approach has been adopted by Gingeras *et al.* (1987) who carried out sandwich hybridization in solution. The target DNA is hybridized in solution to a labelled probe and to an unlabelled capture probe which is immobilized by covalent attachment at its 5' end to a solid support. If the solid support is in the form of beads, subsequent




Fig. 15.12 The sandwich hybridization method as carried out in a microtitre tray. The different wells of the microtitre tray show different stages in the procedure.

